

Be Ready if a Measles Outbreak Affects Your Hospital

Recent measles outbreaks are making headlines.

A common scenario is when an unprotected traveler picks up measles while abroad...then spreads it to unvaccinated people back home.

Nine out of 10 nonimmune patients will get measles after exposure...and about 1 in 4 or 5 will end up in the hospital with complications (pneumonia, etc).

Expect to see airborne isolation precautions for patients admitted with measles symptoms...high fever, runny nose, cough, pink eye, and a red rash that starts on the face.

Keep in mind, measles is highly contagious, especially from 4 days before to 4 days after the rash starts.

Expect these patients to be treated with fluids and meds to control fever (acetaminophen, etc).

Also be ready to dispense oral vitamin A, especially for children...they're likely to be deficient. Measles can decrease vitamin A levels...and supplementing may decrease complications and severity.

Anticipate dispensing the MMR vaccine for any unvaccinated patients that are exposed to measles. Giving it within 3 days of exposure may help prevent measles or limit severity.

But watch for patients who should NOT get MMR. There's concern the live vaccine could lead to complications in patients with a severely weakened immune system...or birth defects in pregnant patients.

Read vaccine orders carefully...and clarify abbreviations. "MMR" can be mistaken for "MMRV" (measles/mumps/rubella/varicella).

If possible, store the MMR vaccine and its diluent together in the fridge. This helps ensure MMR is reconstituted with the correct diluent...and that the diluent isn't injected alone.

Key References:

- CDC. Stay Alert for Measles Cases. January 25, 2024.
<https://emergency.cdc.gov/newsletters/coca/2024/012524.html> (Accessed February 1, 2024).
- CDC. Measles (Rubeola) for Healthcare Providers. November 5, 2020.
<https://www.cdc.gov/measles/hcp/index.html> (Accessed February 1, 2024).
- Kobaidze K, Wallace G. Forgotten but Not Gone: Update on Measles Infection for Hospitalists. J Hosp Med. 2017 Jun;12(6):472-476.
- Wang ME, Ratner AJ. Clinical Guideline Highlights for the Hospitalist: Diagnosis and Management of Measles. J Hosp Med. 2020 Jan 1;15(1):47-48.
- McLean HQ, Fiebelkorn AP, Temte JL, Wallace GS; Centers for Disease Control and Prevention. Prevention of measles, rubella, congenital rubella syndrome, and mumps, 2013: summary recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR Recomm Rep. 2013 Jun 14;62(RR-04):1-34.

Hospital Pharmacy Technician's Letter. March 2024, No. 400328

Cite this document as follows: Article, Be Ready if a Measles Outbreak Affects Your Hospital, Hospital Pharmacist's Letter, March 2024

The content of this article is provided for educational and informational purposes only, and is not a substitute for the advice, opinion or diagnosis of a trained medical professional. If your organization is interested in an enterprise subscription, email sales@trchealthcare.com.

© 2024 Therapeutic Research Center (TRC). TRC and Hospital Pharmacist's Letter and the associated logo(s) are trademarks of Therapeutic Research Center. All Rights Reserved.